



Base Excess and Beyond: Evolving Concepts in Acid-base Analysis

Özge Köner¹, Tuğhan Utku¹, Kubilay Demirağ², Levent Dösemeci³

¹Yeditepe University Faculty of Medicine, Department of Anesthesiology and Intensive Care, İstanbul, Türkiye

²Ege University Faculty of Medicine, Department of Anesthesiology and Intensive Care, İzmir, Türkiye

³Memorial Antalya Hospital, Clinic of Anesthesiology and Intensive Care, Antalya, Türkiye

Abstract

Base excess (BE), a marker used to detect metabolic acid-base disturbances, is also known to predict mortality in critically ill patients; the traditional concept, originally based on the Henderson-Hasselbalch model, has been further refined through integration with the Stewart approach, enabling a more comprehensive and mechanistic evaluation of acid-base disturbances. However, the increasingly complex mathematical formulations required for this integration demand extensive calculations, which can hinder bedside assessment. To address this, the BE formula has been simplified and integrated into the Stewart concept, resulting in a more reliable, detailed, and rapid bedside evaluation. Additionally, the term “alactic BE” was introduced to distinguish metabolic acidosis caused by retention of fixed acids from that caused by lactic acid accumulation, particularly in patients with renal failure. This review discusses the concept of BE and its evolution over the years.

Keywords: Acid-base equilibrium, alactic base excess, base excess, metabolic acidosis, metabolic alkalosis, Stewart approach

Main Points

- The original concept of base excess (BE) was later expanded upon by the Fencl-Stewart model, which reframed the idea in terms of physicochemical principles.
- The Fencl-Stewart equations were rather intricate and impractical for quick calculations. To address this, a simplified variation of the equation that maintains the core physicochemical concepts was introduced.
- The recent concept of “alactic BE” helps clinicians differentiate metabolic acidosis due to lactate accumulation from metabolic acidosis caused by retained fixed acids in patients with renal impairment.
- While the modern BE framework is more effective at identifying mixed acid-base disturbances, its clinical relevance is still uncertain. The recent concept of “alactic BE” helps clinicians differentiate metabolic acidosis due to lactate accumulation from metabolic acidosis caused by retained fixed acids in patients with renal impairment.
- While the modern BE framework is more effective at identifying mixed acid-base disturbances, its clinical relevance is still uncertain.

Introduction

Base excess (BE) is a fundamental parameter used to quantify metabolic acid-base disturbances and has long been recognized for its clinical utility. Beyond its diagnostic role, accumulating evidence indicates that BE also carries prognostic significance in critically ill populations.¹⁻³ In patients with acute kidney injury (AKI), both markedly low and high BE values on admission have been associated with an increased risk of 30-day mortality.¹ Similarly, in critically ill patients with acute myocardial infarction, low BE has been identified as an independent predictor of short- and long-term mortality.² A recent study in patients with ischemic stroke further demonstrated that a BE value below -3 mmol L^{-1} at intensive care unit (ICU) admission may indicate elevated mortality risk.³ Collectively, these findings highlight the potential of BE as a rapid and accessible early risk-stratification tool across a variety of acute and emergency conditions, including AKI, ischemic stroke, cardiac ischemia, and metabolic disturbances. Since its first description in 1948, the concept of BE has undergone substantial refinement and has been incorporated into contemporary approaches to acid-base interpretation, including the modern Stewart

Corresponding author: Prof. Özge Köner MD, PhD, **E-mail:** ozgekoner@gmail.com

Received: June 28, 2025 **Accepted:** December 12, 2025 **Epub:** January 07, 2026 **Published date:** February 09, 2026

Cite this article as: Köner Ö, Utku T, Demirağ K, Dösemeci L. Base excess and beyond: evolving concepts in acid-base analysis. *Turk J Anaesthesiol Reanim.* 2026;54(1):9-14



physicochemical framework. In this review, we examine the conceptual development of BE, its clinical applications, and its evolving role in modern acid-base evaluation.

Buffer Base Concept

Buffer base (BB), which was introduced by Singer and Hastings⁴ in 1948, is the predecessor of the BE concept. According to this concept, plasma BB is practically equal to the sum of all buffer anions [total base (bicarbonate) and weak acid anions (e.g., albumin and phosphate)]. Therefore, BB increased either by an increase in base or by a decrease in plasma albumin concentration (Equation 1).

$$BB = HCO_3^- + A^- \quad (1)$$

The BB concept considers the non-carbonic buffers and is theoretically CO_2 -independent. Unfortunately, inter-subject variability due to differences in non-carbonic buffer concentrations was observed. To overcome this limitation, Siggaard-Andersen introduced the BE concept, in which the “excess” may be positive or negative, and which defines the actual BB relative to the normal BB (i.e., the actual BB minus the BB at normal pH and pCO_2) as a measure of metabolic acid-base disturbance.^{5,6}

Base Excess Concept

BE is the amount of acid or base ($mmol\ L^{-1}$) that must be added to a blood sample to reach a pH of 7.40 in under standardized conditions ($pCO_2\ 40\ mmHg$, $37\ ^\circ C$).⁶ Affecting factors affecting of this semi-quantitative approach are pCO_2 and BB, and; standard BE is used as its marker. BE is a Henderson Hasselbalch based Henderson-Hasselbalch-based parameter. It is calculated from the Van Slyke equation, which accounts for pH, HCO_3^- ($mmol\ L^{-1}$) and haemoglobin (Hb, $mmol\ L^{-1}$) (Equation 2).⁷

$$BE = (HCO_3^- - 24.4 + [(2.3 \times Hb + 7.7) \times (pH - 7.40)]) + (1 - 0.023 \times Hb) \quad (2)$$

In this BE formula of Van Slyke, Hb can either be a constant value ($16.2\ mmol\ L^{-1}$) or be computed as a function of Hb concentration (assuming a constant protein concentration of $70\ g\ L^{-1}$). The BE formula has also exhibits some inconsistencies due to changes in CO_2 . As in vivo BE is less reliable than in vitro BE, Van Slyke modified the equation to standardize the effect of Hb on BE [standard base excess (SBE)] and considered a lower Hb value for the calculation (Equation 3).

$$SBE = 0.9287 [HCO_3^- - 24.4 + 14.83 \times (pH - 7.40)] \quad (3)$$

However, the SBE equation assumes that the plasma albumin ($g\ dL^{-1}$) and phosphate ($mg\ dL^{-1}$) concentrations (Atot) are within the normal range. Therefore, Wooten described another formula for a multi-compartment model (Equation 4).^{8,9} The normal value of SBE is between -2 and $+2\ mmol\ L^{-1}$.

$$\text{Corrected SBE} = (HCO_3^- - 24.4) + [(8.3 \times \text{albumin} + 0.15) + (0.29 \times \text{Pi} \times 0.32) \times (pH - 7.40)] \quad (4)$$


This multi-compartmental model has been shown to correlate well with the strong ion difference (SID) concept described by Stewart in that the change in SBE equals the change in SID across a vascular bed if there is no change in Atot.^{8,10,11} The anionic contribution of the most important weak acid, albumin (Atot), is $10\ mEq\ L^{-1}$, whereas, unless phosphate levels are increased, their contribution to Atot is around 5%. If phosphate levels are elevated, then its contribution becomes more important. Because it is difficult to calculate Stewart's SID equation, BE can be used as a substitute for SID at the bedside. Standard BE is a powerful concept in defining the metabolic acid-base disorders; however, it is not helpful for diagnosing the underlying pathology.¹² Therefore, “mixed” acid-base disorders will go unnoticed without the use of detailed base-excess partitioning.^{13,14} Furthermore, in some pathological situations, BE may be within the normal range and thus be misleading (Figure 1).¹⁵

Gamblegram^{8,16} is a graphic representation of SID (therefore BE) developed by James Gamble (Figure 1). Assuming electroneutrality, the Gamblegram demonstrates that the ions that occupy the SID between the strong cations and anions are primarily bicarbonate (a component dependent on the SID) and the total concentration of weak acids.

Ciabattoni et al.¹⁶ compared Stewart's SID and SBE approaches for detecting metabolic disturbances in a group of ICU patients receiving mechanical ventilation and found that the correlation between the two parameters was poor. They explained the results by stating that the SBE was derived from a variety of parameters, showed the effect of a single independent variable in the Stewart approach.¹⁶ Stewart's approach was superior to SBE in identifying metabolic alkalosis related to hypoalbuminemia and hypophosphatemia, and in identifying metabolic acidosis related to haemodilution following fluid resuscitation.

Stewart Approach and its integration to BE concept

Unlike the classical Henderson-Hasselbalch approach, Stewart demonstrated that the concentration of $[H^+]$ in a physiological fluid is determined by three independent variables: the SID (which is equal to the difference between positively and negatively charged ions that are fully dissociated in biological fluids, e.g., Na^+ , K^+ , Cl^- , lactate), the total concentration of weak acids (Atot), and the volatile acid CO_2 . According to Stewart, $[H^+]$ and $[HCO_3^-]$ are dependent variables, and bicarbonate appears to be an indicator rather than a cause. Water becomes ionized under the influence of these independent variables, thereby regulating the pH. Stewart conducted his acid-base studies by applying the principles of electrical neutrality, the law of mass action, and conservation of mass. When expressed



Figures 1 A, B. Gamblegrams show the standard base excess values of two patients, both are within the normal range (0). Figure 1 A represents a patient with normal acid-base balance, whereas Figure 1 B represents a patient with hyperlactatemia receiving a bicarbonate infusion. SBE, standard base excess; A-, non-volatile weak acids.¹²

mathematically, apparent SID (SIDa) is equivalent to the “plasma BB” described by Singer and Hastings.⁴

Fencl-Stewart Approach

To overcome the weaknesses of the SBE concept, Balasubramanyan et al.¹⁷ derived a new formula from the work of Fencl and defined it as the Fencl-Stewart approach. This derived formula combines BE with the Stewart approach. Gilfix et al.¹⁸ derived an equation to estimate the BE effects of the SID and the Atot. However, this formula is complex and difficult to apply at the bedside (Equation 5). Therefore, another equation, a simplified form of the Fencl-Stewart formula, was described by Story et al.¹⁹ (Equation 6).

$$BE = 0.3x [(\text{Na}^+) - 140] + 102 - [(\text{Cl}^-) \times 140 / (\text{Na}^+)] + (0.123x \text{pH} - 0.631) \times [42 - \text{albumin (g L}^{-1})] \quad (5)$$

Story's Simplified Fencl-Stewart Approach

Rather than the complex formula of the aforementioned Fencl-Stewart approach, Story et al.¹⁹ proposed a simplified form (Equation 6). They confirmed in a study of 300 blood samples taken from critically ill patients that this simplified formula was in good agreement with the more complex Fencl-Stewart approach. Using this formula, the effects of SID and albumin on BE can be estimated.

$$\text{SBE NaCl} = \{[\text{Na}^+] - [\text{Cl}^-]\} - 38 \quad (38 \text{ is the average normal SID}) \quad (6)$$

$$\text{SBE Alb} = 0.25 \times (42 - \text{measured albumin in blood}) \quad (42 \text{ is the normal plasma albumin g L}^{-1})$$

$$\text{SBE NaCl} + \text{SBE Alb} = \text{SBE Correction}$$

$$\text{True SBE or BE gap} = \text{SBE} - \text{SBE Correction}$$

The above equations require only basic mental arithmetic and are therefore easy to use at the bedside for evaluation, treatment, and follow-up of both simple and complex acid-base problems.

Alactic Base Excess

Alactic BE (ABE) was defined by Gattinoni et al.²⁰ to differentiate metabolic acidosis related to retention of non-lactate fixed acids (sulphates, phosphates) from that related to lactic acid in patients with renal failure (Equation 7).

$$\text{ABE (mmol L}^{-1}) = \text{SBE (mmol L}^{-1}) + \text{Lactate (mmol L}^{-1}) \quad (7)$$

Clinical and Research Consequences

Studies with Fencl-Stewart approach

Traditional BE and bicarbonate methods can miss complex metabolic disorders in critically ill patients. While the

Stewart and modified BE methods better identify multiple coexisting disturbances, their routine use may be more labor-intensive. The modified BE approach offers a practical balance between ease of use and clinical accuracy.²¹ In liver transplant patients, the Fencl-Stewart approach helps identify complex acid-base changes immediately after transplantation.²² Fencl-Stewart's method revealed frequent simultaneous metabolic acidosis and alkalosis missed by traditional analysis; both strong ion gap (SIG) and lactate independently predicted 28-day ICU mortality in critically ill.²³

Balasubramanyan et al.¹⁷ found that unmeasured anions measure with Fencl-Stewart's BE method correlates with mortality in pediatric intensive care unit patients, whereas Cusack et al.²⁴ did not observe this in adults. To address the small sample sizes of earlier studies, Rocktaeschel et al.²⁵ retrospectively analyzed 300 adult intensive care unit patients and found that unmeasured anions were the only acid-base variables with limited ability to predict mortality, while all calculation methods [anion gap (AG), corrected AG, SIG, and unmeasured anions measured by Fencl-Stewart BE] were strongly correlated with each other. Furthermore, of four methods for assessing chloride's effect on acid-base status, chloride-specific base excess (BECl) most closely correlates with standard BE and most accurately reflects its impact in critically ill patients.²⁶

A clinical example of simplified Fencl-Stewart approach by Story et al.¹⁹

The patient underwent major gynaecological surgery with general anaesthesia. Isotonic sodium chloride was used as an intraoperative fluid. The metabolic acidosis, which developed 2 hours after induction of anaesthesia, is due to a decreased SID related to hyperchloremia. This effect was offset by a decrease in the concentration of albumin (a weak acid). There is no contribution from unmeasured ions in this case. This acidosis was observed after the infusion of 6 litres of isotonic sodium chloride (Table 1).

Table 1. Simplified Fencl-Stewart Approach¹⁹

| Measured values | After anaesthesia induction | 2 hours after anaesthesia induction |
|--|-----------------------------|-------------------------------------|
| pH | 7.41 | 7.28 |
| CO ₂ (mmHg) | 39.7 | 39.7 |
| Na ⁺ | 140 | 142 |
| Cl ⁻ | 104 | 115 |
| Base excess (mEq L ⁻¹) | -0.4 | -6.7 |
| Albumin (g dL ⁻¹) | 4 | 2.8 |
| Na-Cl effect (mEq L ⁻¹) | -2 | -11 |
| Albumin effect (mEq L ⁻¹) | 0.5 | 3.5 |
| Effect of unmeasured anions (mEq L ⁻¹) | 1.1 | 0.8 |

$$\text{Na} - \text{Cl effect on BE} = (\text{Na}^+) - (\text{Cl}^-) - 38$$

$$\text{Albumin effect on BE} = 0.25x [42 - (\text{Albumin}) - \text{g L}^{-1}]$$

$$\text{Unmeasured ion effect on BE} = \text{SBE} - (\text{Na}^+ - \text{Cl}^-) - \text{Albumin effect}$$

A clinical example comparing Fencl-Stewart, anion gap, and Stewart methods.²¹

Table 2 presents a patient with postoperative multiple organ failure. The patient has complex acid-base physiology. The pH of 7.33, pCO₂ of 30, and BE (-10) of -10 suggest metabolic acidosis and respiratory alkalosis; however, the etiology of the metabolic acidosis cannot be identified. By the AG method, the elevated corrected AG ($\Delta\text{AGcorr} = 11 \text{ mEq L}^{-1}$) suggests metabolic acidosis due to unmeasured anions. The delta-delta calculation predicts a bicarbonate level of 13-17 mEq L⁻¹, and the observed bicarbonate of 15 mEq L⁻¹ falls within this expected range. Based on this approach, no additional metabolic acid-base disorders are identified, even though a hypoalbuminemic alkalosis is intuitively suspected. In contrast, the Stewart approach reveals several abnormalities: an increased SIG, reduced sodium levels, elevated corrected chloride levels, and decreased albumin and phosphate levels. Together, these findings indicate a complex disturbance consisting of multiple metabolic acidoses (due to unmeasured anions, free-water excess, and hyperchloremia) and concurrent hypoalbuminemic and hypophosphatemic alkalosis. The modified BE method yields similar detail: a positive BEalb (+13 mEq L⁻¹), a negative BECl (-8 mEq L⁻¹), a negative BE for free water, and a negative BE for unmeasured anions (-8 mEq L⁻¹), reflecting coexisting hypoalbuminemic alkalosis, hyperchloremic acidosis, dilutional acidosis, and unmeasured-anion acidosis. In this case, only the Stewart and modified BE methods accurately characterize all metabolic acid-base disturbances, whereas the AG method fails to detect the hypoalbuminemic alkalosis, hyperchloremic acidosis, and dilutional acidosis. The clinical significance of these missed abnormalities, however, remains uncertain.

Table 2. Comparing Fencl-Stewart, Anion Gap, and Stewart Methods²¹

| Measured values | Data |
|--|------|
| pH | 7.33 |
| CO ₂ (mmHg) | 30 |
| Na ⁺ (mEq L ⁻¹) | 117 |
| Cl ⁻ (mEq L ⁻¹) | 104 |
| Cl corrected (mEq L ⁻¹) | 112 |
| K ⁺ (mEq L ⁻¹) | 3.9 |
| BE lab (mEq L ⁻¹) | -10 |
| Albumin (g dL ⁻¹) | 6 |
| Pi (mmol L ⁻¹) | 0.6 |
| HCO ₃ ⁻ | 15 |
| ΔAgap corrected | 11 |
| Strong ion gap | 18 |

Alactic base excess

Gattinoni et al.²⁰ studied ABE in septic patients and suggested that ABE may reflect the effects of renal dysfunction on plasma lactate concentration. Negative ABE values ($< -3 \text{ mmol L}^{-1}$) are correlated with mortality in patients with sepsis. Neutral ABE values ($\geq -3 \text{ to } < 4 \text{ mmol L}^{-1}$) indicate that the kidneys effectively excrete fixed acids and maintain physiologic blood pH. Positive ABE values ($\geq 4 \text{ mmol L}^{-1}$) suggest that lactate and standard base levels are within the normal range in the blood, reflecting the kidney's compensatory response to metabolic acidosis. In their study of intensive care patients with shock, Smuszkiewicz et al.²⁷ reported that an admission ABE below $-3.63 \text{ mmol L}^{-1}$ may be associated with an increased risk of 28-day all-cause mortality. Moreover, both BE and lactate levels—each contributing to ABE—offer independent and complementary predictive value.²⁷ The combined criteria of BE $< -9.5 \text{ mmol L}^{-1}$ and lactate $> 4.5 \text{ mmol L}^{-1}$ identify patients at the greatest risk of death more effectively than BE, lactate, or ABE alone. Importantly, the prognostic significance of these metabolic markers remains robust regardless of patients' age, sex, type of shock, or presence of severe renal failure. Negative ABE is also an independent predictor of in-hospital mortality among septic patients both with and without renal dysfunction, and among patients with acute myocardial infarction.^{28,29}

Conclusion

In the mid-20th century, the concept of BB, which represents the sum of the concentrations of bicarbonate, albumin, and haemoglobin, evolved into the concept of delta BB, defined as the change in BB from the “normal” value. Since the blood volume is diluted by interstitial fluid, the extracellular fluid model was introduced; this model uses a standard haemoglobin concentration (5 g dL^{-1} or 3 mmol L^{-1}). Although it reflects only the non-respiratory (metabolic) component of acid-base disorders, the BE of the extracellular fluid remains valuable for etiological analysis, particularly when used alongside the Stewart approach in partitioning contributing factors. Clearly, the concept of BE continues to evolve, and ongoing refinements are likely. The Fencl-Stewart concept of BE has now been refined using the formula proposed by Story et al.¹⁹ This enables straightforward bedside calculations and helps identify the underlying pathology of metabolic acid-base disturbances, which the BE value alone cannot provide. The novel ABE concept further enhances our understanding by shedding light on the kidney's role in metabolic acidosis.

Footnotes

Author Contributions: Concept – Ö.K., T.U., K.D., L.D.; Design - Ö.K., T.U., K.D., L.D.; Data Collection or Processing - Ö.K., T.U., K.D., L.D.; Analysis or Interpretation - Ö.K., T.U., K.D., L.D.;

Literature Review - Ö.K., T.U.; Writing - Ö.K., T.U.

Declaration of Interests: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding: The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

- Cheng Y, Zhang Y, Tu B, et al. Association between base excess and mortality among patients in icu with acute kidney injury. *Front Med (Lausanne)*. 2021;8:779627. [\[CrossRef\]](#)
- Luo C, Duan Z, Zheng T, et al. Base excess is associated with the risk of all-cause mortality in critically ill patients with acute myocardial infarction. *Front Cardiovasc Med*. 2022;9:942485. [\[CrossRef\]](#)
- Liu J, Li J, Jin X, et al. Association between base excess and mortality in critically ill patients with ischemic stroke: a retrospective cohort study. *BMC Neurol*. 2024;24(1):351. [\[CrossRef\]](#)
- Singer RB, Hastings AB. An improved clinical method for the estimation of disturbances of the acid-base balance of human blood. *Medicine (Baltimore)*. 1948;27(2):223-242. [\[CrossRef\]](#)
- Astrup P, Jorgensen K, Andersen OS, Engel K. The acid-base metabolism. A new approach. *Lancet*. 1960;1(7133):1035-1039. [\[CrossRef\]](#)
- Siggaard-Andersen O. The acid-base status of the blood. *Scand J Clin Lab Invest*. 1963;15 Suppl 70:1-134. [\[CrossRef\]](#)
- Van Slyke DD. An apparatus for determination of the gases in blood and other solutions. *Proc Natl Acad Sci U S A*. 1921;7(8):229-231. [\[CrossRef\]](#)
- Wooten EW. Calculation of physiological acid-base parameters in multicompartment systems with application to human blood. *J Appl Physiol (1985)*. 2003;95(6):2333-2344. [\[CrossRef\]](#)
- Wooten EW. Science review: quantitative acid-base physiology using the Stewart model. *Crit Care*. 2004;8(6):448-452. [\[CrossRef\]](#)
- Schllichtig R. [Base excess] vs [strong ion difference]. Which is more helpful? *Adv Exp Med Biol*. 1997;411:91-95. [\[CrossRef\]](#)
- Wooten EW. Analytic calculation of physiological acid-base parameters in plasma. *J Appl Physiol (1985)*. 1999;86(1):326-334. [\[CrossRef\]](#)
- Tuhay G, Pein MC, Masevicius FD, Kutscherauer DO, Dubin A. Severe hyperlactatemia with normal base excess: a quantitative analysis using conventional and Stewart approaches. *Crit Care*. 2008;12(3):R66. [\[CrossRef\]](#)
- Funk GC, Doberer D, Osterreicher C, Peck-Radosavljevic M, Schmid M, Schneeweiss B. Equilibrium of acidifying and alkalinizing metabolic acid-base disorders in cirrhosis. *Liver Int*. 2005;25(3):505-512. [\[CrossRef\]](#)
- Ahya SN, José Soler M, Levitsky J, Batlle D. Acid-base and potassium disorders in liver disease. *Semin Nephrol*. 2006;26(6):466-470. [\[CrossRef\]](#)
- Langer T, Brusatori S, Gattinoni L. Understanding base excess (BE): merits and pitfalls. *Intensive Care Med*. 2022;48(8):1080-1083. [\[CrossRef\]](#)
- Ciabattoni A, Chiumello D, Mancusi S, et al. Acid-base status in critically ill patients: physicochemical vs. traditional approach. *J Clin Med*. 2025;14(9):3227. [\[CrossRef\]](#)
- Balasubramanyan N, Havens PL, Hoffman GM. Unmeasured anions identified by the Fencl-Stewart method predict mortality better than base excess, anion gap, and lactate in patients in the pediatric intensive care unit. *Crit Care Med*. 1999;27(8):1577-1581. [\[CrossRef\]](#)
- Gilfix BM, Bique M, Magder S. A physical chemical approach to the analysis of acid-base balance in the clinical setting. *J Crit Care*. 1993;8(4):187-197. [\[CrossRef\]](#)

19. Story DA, Morimatsu H, Bellomo R. Strong ions, weak acids and base excess: a simplified Fencl-Stewart approach to clinical acid-base disorders. *Br J Anaesth.* 2004;92(1):54-60. [\[CrossRef\]](#)
20. Gattinoni L, Vasques F, Camporota L, et al. Understanding lactatemia in human sepsis. Potential impact for early management. *Am J Respir Crit Care Med.* 2019;200(5):582-589. [\[CrossRef\]](#)
21. Fidkowski C, Helstrom J. Diagnosing metabolic acidosis in the critically ill: bridging the anion gap, Stewart, and base excess methods. *Can J Anaesth.* 2009;56(3):247-256. [\[CrossRef\]](#)
22. Story DA, Vaja R, Poustie SJ, McNicol L. Fencl-Stewart analysis of acid-base changes immediately after liver transplantation. *Crit Care Resusc.* 2008;10(1):23. [\[CrossRef\]](#)
23. Antonini B, Piva S, Paltenghi M, Candiani A, Latronico N. The early phase of critical illness is a progressive acidic state due to unmeasured anions. *Eur J Anaesthesiol.* 2008;25(7):566-571. [\[CrossRef\]](#)
24. Cusack RJ, Rhodes A, Lochhead P, et al. The strong ion gap does not have prognostic value in critically ill patients in a mixed medical/surgical adult ICU. *Intensive Care Med.* 2002;28(7):864-869. [\[CrossRef\]](#)
25. Rocktaeschel J, Morimatsu H, Uchino S, Bellomo R. Unmeasured anions in critically ill patients: can they predict mortality? *Crit Care Med.* 2003;31(8):2131-2136. [\[CrossRef\]](#)
26. Gucyetmez B, Tuzuner F, Atalan HK, Sezerman U, Gucyetmez K, Telci L. Base-excess chloride; the best approach to evaluate the effect of chloride on the acid-base status: a retrospective study. *PLoS One.* 2021;16(4):e0250274. [\[CrossRef\]](#)
27. Smuszkiewicz P, Jawień N, Szrama J, Lubarska M, Kusza K, Guzik P. Admission lactate concentration, base excess, and alactic base excess predict the 28-day inward mortality in shock patients. *J Clin Med.* 2022;11(20):6125. [\[CrossRef\]](#)
28. Cantos J, Huespe IA, Sinner JF, et al. lactic base excess is an independent predictor of death in sepsis: a propensity score analysis. *J Crit Care.* 2023;74:154248. [\[CrossRef\]](#)
29. Zhou C, Wang Q. Association of alactic base excess with in-hospital mortality in patients with acute myocardial infarction: a retrospective cohort study. *BMC Cardiovasc Disord.* 2024;24(1):447. [\[CrossRef\]](#)